

# PROFESSOR. IAN T. JONES

FRACS FRCS

## COLORECTAL SURGERY

Suite 5, 6 & 7  
Private Medical Centre  
Royal Melbourne Hospital Royal Parade  
Parkville VIC 3050

Tel: 9347 4077  
Fax: 9348 2075  
Pager: 9387 1000  
www.colorectalmelbourne.com.au

---

## COLONOSCOPY PREPARATION (COLONLYTELY)

A colonoscopy is a visual examination of the lining of your colon (large intestine). A flexible tube (colonoscope) is passed around the bowel and a careful search made for any abnormalities. If necessary, small tissue samples (biopsies) can be taken or abnormal growths (polyps) can be removed during the procedure. Intravenous medication is given by an anaesthetist during the procedure so patient discomfort is minimal. You will be informed where the colonoscopy will be undertaken and the time to attend. Please note this is the time to arrive which is usually one to two hours before the procedure.

### PREPARATION

To allow a clear view, the colon must be free of waste material. You must drink a special cleansing solution on the afternoon of the day before the procedure. The recommended product is COLONLYTELY. This is available from your chemist WITHOUT PRESCRIPTION.

## THE DAY BEFORE YOUR COLONOSCOPY

- No Solids can be eaten on this day. DRINK ONLY CLEAR LIQUIDS. (Thick liquids or food must not be consumed. See over for list of permitted clear fluids.)
- Mix each sachet in a separate litre of water for a total volume of THREE (3) litres
- This needs to be drunk at the rate of ONE LITRE PER HOUR in between the times of 2PM and 6PM on the day before the colonoscopy
- In addition, drink clear liquids at least one glass per hr (black teas & coffee, lemonade, jelly (not red), clear apple juice

## THE DAY OF THE COLONOSCOPY

For a **MORNING** colonoscopy:

- Stop all oral intake from midnight until after your colonoscopy

For an **AFTERNOON** colonoscopy:

- Stop all oral intake from 8 a.m. on the day of the colonoscopy

## AVOIDING DEHYDRATION

As mentioned above, once you start the bowel preparation you are restricted to clear liquids until the specified fasting time. This does not mean that you need to restrict the amount of liquids you drink. Indeed you are encouraged to take an adequate amount of fluid as failure in sufficient quantity can leave you dehydrated.

## APPROVED CLEAR FLUIDS

- Water
- Clear broths
- Black tea or black coffee
- Sports drinks eg Gatorade
- Carbonated beverages
- Barley sugar

## MEDICATIONS

Even though you will undertake a period of fasting before the colonoscopy, you should take your usual prescribed medications at the usual time with a SIP of water. **SOME MEDICATIONS SHOULD BE STOPPED BEFORE THE COLONOSCOPY.** These include WARFARIN, ASPIRIN, CLOPIDOGREL AND IRON. ASPIRIN products should be avoided for SEVEN DAYS and IRON PREPARATIONS for FOUR DAYS before the procedure.

**PLEASE DISCUSS THIS WITH PROFESSOR JONES.**

Please also advise us if you have had rheumatic fever or heart valve surgery.

## AFTER THE COLONOSCOPY

You will remain under observation for 1-½ hours until the effect of the sedation medication wears off. Because of the medication you must NOT drive a car, operate machinery, make important decisions or drink alcohol for the rest of the day. **IT IS ESSENTIAL FOR A FRIEND OR RELATIVE TO ACCOMPANY YOU HOME.**

## RISKS

Colonoscopy can result in complications such as damage to or bleeding from the bowel. These complications are rare (less than 1 in 1000) but may require urgent treatment or even major surgery to correct. There is a remote risk of death (less than 1 in 150,000 cases following colonoscopy)

## ADMISSION TIME & PLACE

HOSPITAL: \_\_\_\_\_

DATE: \_\_\_\_\_

TIME: \_\_\_\_\_ AM / PM

Ian T. Jones